

Supplementary Figures: S2A-F

AcetoBase Version 2: A database update and re-analysis of formyltetrahydrofolate synthetase amplicon sequencing data from anaerobic digesters

Abhijeet Singh^{1*}, Anna Schnürer^{1*}

¹Anaerobic Microbiology and Biotechnology Group, Department of Molecular Sciences, Swedish University of Agricultural Sciences, Almas Allé 5, Uppsala, SE-750 07, Uppsala, Sweden

*For correspondence. E-mail: anna.schnurer@slu.se; abhijeet.singh@slu.se, Tel. +46 18671000; Fax +46 18672000; Address: Department of Molecular Sciences, Box 7025, 75007 Uppsala, Sweden

Description:


The supplementary figures S2A-F (phylum-species level) are generated in present study by the reanalysis of the FTHFS amplicon sequence data from the study by Singh *et al.* (2021).

Reference:

Singh, A., Müller, B., and Schnürer, A. (2021). Profiling temporal dynamics of acetogenic communities in anaerobic digesters using next-generation sequencing and T-RFLP. *Sci. Rep.* 11, 13298. doi:10.1038/s41598-021-92658-2.


S2A

FTHFS-community: Phylum Relative Abundance (> 1 %)




S2B


FTHFS-community: Class Relative Abundance (> 1 %)



S2C

FTHFS-community: Order Relative Abundance (> 1 %)





FTHFS-community: Genus Relative Abundance (> 1 %)

S2E

