

Guidelines on the diagnosis and management of acute pulmonary embolism

The Task Force for the Diagnosis and Management of Acute Pulmonary Embolism of the European Society of Cardiology (ESC)

Supplementary material

Table A Major trials reporting RV/LV diameter ratio assessed by CT as a risk marker for 30-day all-cause mortality in acute PE

Author	n	CT equipment	Cutoff	Positive (%)	Sensitivity (%)	Specificity (%)	NPV (%)	PPV (%)
Van der Meer <i>et al.</i> ¹	120	SDCT	RV/LV >1	57.5	NA	NA	100	10
Schoepf <i>et al.</i> ²	431	4–16 MDCT	RV/LV >0.9	64	78.2	38	92.3	15.6

n = number of patients; SDCT = single-detector computed tomography; MDCT = multidetector computed tomography; RV/LV = ratio of diastolic diameters of RV and LV; NA = not available; NPV = negative predictive value; PPV = positive predictive value for all-cause 30-day mortality.

Table B Major trials on the prognostic value of BNP/NT-proBNP in acute pulmonary embolism

Author	n	Biomarker and test used	Threshold	Positive (%)	Sensitivity (%)	Specificity (%)	NPV (%)	PPV ^a (%)
Ten Wolde <i>et al.</i> ³	110	BNP ^b	21.7 pmol/L	33	86	73	99	18
Kucher <i>et al.</i> ⁴	73	NT-proBNP ^c	500 pg/mL	58	NA	NA	100	12
Kucher <i>et al.</i> ⁵	73	BNP ^d Triage	50 pg/mL	58	NA	NA	100	12
Pruszczyk <i>et al.</i> ⁶	79	NT-proBNP ^c	600 pg/mL	27	100	33	100	26
Binder <i>et al.</i> ⁷	124	NT-proBNP ^c	1000 pg/mL	54	100	49	100	10
Kostrubiec <i>et al.</i> ⁸	113	NT-proBNP ^c	NT-proBNP >7500 pg/mL ^a	16	65	93	94	61

^aIf not decreasing by 50% within 24 h

^{b–d}Tests used: ^bShionoria, CIS Bio International; ^cElecsys, Roche Diagnostics; ^dTriage, Biosite Technologies.

n = number of patients; BNP = brain natriuretic peptide; NT-proBNP = N-terminal proBNP; NA = not available; NPV = negative predictive value; PPV = positive predictive value.

Table C Major studies on the prognostic value of biochemical markers of myocardial injury in pulmonary embolism

Author	n	Biomarker and test used	Threshold	Positive (%)	Sensitivity (%)	Specificity (%)	NPV (%)	PPV (%)
Troponins								
Giannitsis <i>et al.</i> ⁹	56	cTnT ^a	0.10 ng/mL	32	89	79	97	44
Konstantinides <i>et al.</i> ¹⁰	106	cTnI ^b	0.07 ng/mL	41	86	63	98	14
Konstantinides <i>et al.</i> ¹⁰	106	cTnT ^a	0.04 ng/mL	37	71	66	97	12
Janata <i>et al.</i> ¹¹	106	cTnT ^a	0.09 ng/mL	11	80	92	99	34
Pruszczyk <i>et al.</i> ¹¹	64	cTnT ^a	0.01 ng/mL	50	100	57	100	25
Douketis <i>et al.</i> ¹³	458	cTnI ^c	0.5 ng/mL	13.5	For all-cause 90-day mortality: odds ratio 3.5, 95% CI 1.0–11.9			
Other markers of injury								
Pruszczyk <i>et al.</i> ¹⁴	46	Myoglobin ^a	58 ng/mL (women); 72 ng/ml (men)	46	100	64	100	33
Kaczynska <i>et al.</i> ¹⁵	77	H-FABP ^a	6 ng/mL	39	Hazard risk 1.03, 95% CI 1.01–1.05, <i>P</i> < 0.0001		78	23
Puls <i>et al.</i> ¹⁶	107	H-FABP ^d	6 ng/mL	27	100	83	100	37

If not specified otherwise, the data refer to in-hospital mortality.

n = number of patients; cTnI = cardiac troponin I; cTnT = cardiac troponin T; H-FABP = heart-type fatty acid binding protein; NA = not available; NPV = negative predictive value; PPV = positive predictive value.

^{a–d}Tests used: ^aElecsys, Roche Diagnostics, Mannheim, Germany; ^bCentaur, Bayer, Munich Germany; ^cAxSYM, Abbot, Abbott Park, IL, USA; ^dHyCult Biotechnology, Uden, Netherlands.

Table D Prognostic value of concomitant assessment of NT-proBNP and troponin in acute PE

Author	n	Biomarker/ assay used	Threshold	Positive (%)	Endpoint	Sensitivity (%)	Specificity (%)	NPV (%)	PPV (%)
Kostrubiec <i>et al.</i> ¹⁷	100 with systolic blood pressure >90 mmHg	NT-proBNP ^a , cTnT ^a	NT-proBNP <600 pg/mL and cTnT <0.07 µ/L	28	PE-related 40-day mortality	0 ^b	70	89	0
			NT-proBNP >600 pg/mL and cTnT <0.07 µ/L	54	PE-related 40-day mortality	25	44	87	4
			NT-proBNP >600 pg/mL and cTnT >0.07 µ/L	18	PE-related 40-day mortality	75	87	98	33

^aTest used: Elecsys, Roche Diagnostics; Mannheim, Germany.

^bNo death in this group.

n = number of patients; NT-proBNP = N-terminal proBNP; cTnT = cardiac troponin T; NPV = negative predictive value; PPV = positive predictive value.

Table E Prognostic value of concomitant assessment of echocardiography and troponin measurements in PE

Author	n	Troponin assay	Threshold	Echocardiography	Endpoint	Both tests positive (% of patients)	Parameter
Kucher et al. ¹⁸	91	cTnI ^a	>0.06 ng/mL	At least moderate RVD	In-hospital mortality, catecholamines i.v., thrombolysis, CPR, intubation, embolectomy	26.3	Sensitivity 86% Specificity 91% NPV 96% PPV 75%
Binder et al. ⁷	124	cTnT ^b	≥0.04 ng/mL	RV >30 mm Parasternal view	In-hospital mortality, catecholamines iv, thrombolysis, CPR, intubation,	12.9	OR 10.00 (95% CI 2.1–46.8)
Scridon et al. ¹⁹	141	cTnI ^c	≥0.1 ng/mL	RV/LV >0.9	All-cause 30-day mortality	32	OR 7.17 (95% CI 1.6–31.9)

^{a–c}Tests used: ^aAxSYM, Abbott, Abbott Park, IL, USA; ^bElecsys, Roche Diagnostics, Mannheim, Germany; ^cBaxter, Miami, FL, USA.

CI = confidence interval; CPR = cardiopulmonary resuscitation; cTnI = cardiac troponin I; cTnT = cardiac troponin T; LV = left ventricle; n = number of patients; OR = odds ratio; RV = right ventricle; RVD = right ventricular dysfunction.

References

- van der Meer RW, Pattynama PM, van Strijen MJ, van den Berg-Huisman AA, Hartmann JJ, Putter H et al. Right ventricular dysfunction and pulmonary obstruction index at helical CT: prediction of clinical outcome during 3-month follow-up in patients with acute pulmonary embolism. *Radiology* 2005;**235**:798–803.
- Schoepf UJ, Kucher N, Kipfmüller F, Quiroz R, Costello P, Goldhaber SZ. Right ventricular enlargement on chest computed tomography: a predictor of early death in acute pulmonary embolism. *Circulation* 2004;**110**:3276–3280.
- Ten Wolde M, Tulevski II, Mulder JW, Sohne M, Boomsma F, Mulder BJ et al. Brain natriuretic peptide as a predictor of adverse outcome in patients with pulmonary embolism. *Circulation* 2003;**107**:2082–2084.
- Kucher N, Printzen G, Doernhoefer T, Windecker S, Meier B, Hess OM. Low pro-brain natriuretic peptide levels predict benign clinical outcome in acute pulmonary embolism. *Circulation* 2003;**107**:1576–1578.
- Kucher N, Printzen G, Goldhaber SZ. Prognostic role of brain natriuretic peptide in acute pulmonary embolism. *Circulation* 2003;**107**:2545–2547.
- Pruszczyk P, Kostrubiec M, Bochowicz A, Styczynski G, Szulc M, Kurzyna M et al. N-terminal pro-brain natriuretic peptide in patients with acute pulmonary embolism. *Eur Respir J* 2003;**22**:649–653.
- Binder L, Pieske B, Olschewski M, Geibel A, Klostermann B, Reiner C et al. N-terminal pro-brain natriuretic peptide or troponin testing followed by echocardiography for risk stratification of acute pulmonary embolism. *Circulation* 2005;**112**:1573–1579.
- Kostrubiec M, Pruszczyk P, Kaczynska A, Kucher N. Persistent NT-proBNP elevation in acute pulmonary embolism predicts early death. *Clin Chim Acta* 2007;**382**:124–128.
- Giannitsis E, Muller-Bardorff M, Kurowski V, Weidtmann B, Wiegand U, Kampmann M et al. Independent prognostic value of cardiac troponin T in patients with confirmed pulmonary embolism. *Circulation* 2000;**102**:211–217.
- Konstantinides S, Geibel A, Olschewski M, Kasper W, Hruska N, Jackle S et al. Importance of cardiac troponins I and T in risk stratification of patients with acute pulmonary embolism. *Circulation* 2002;**106**:1263–1268.
- Janata K, Holzer M, Laggner AN, Mullner M. Cardiac troponin T in the severity assessment of patients with pulmonary embolism: cohort study. *Br Med J* 2003;**326**:312–313.
- Pruszczyk P, Bochowicz A, Torbicki A, Szulc M, Kurzyna M, Fijalkowska A et al. Cardiac troponin T monitoring identifies high-risk group of normotensive patients with acute pulmonary embolism. *Chest* 2003;**123**:1947–1952.
- Douketis JD, Crowther MA, Stanton EB, Ginsberg JS. Elevated cardiac troponin levels in patients with submassive pulmonary embolism. *Arch Intern Med* 2002;**162**:79–81.
- Pruszczyk P, Bochowicz A, Kostrubiec M, Torbicki A, Szulc M, Gurba H et al. Myoglobin stratifies short-term risk in acute major pulmonary embolism. *Clin Chim Acta* 2003;**338**:53–56.
- Kaczynska A, Pelsers MM, Bochowicz A, Kostrubiec M, Glatz JF, Pruszczyk P. Plasma heart-type fatty acid binding protein is superior to troponin and myoglobin for rapid risk stratification in acute pulmonary embolism. *Clin Chim Acta* 2006;**371**:117–123.
- Puls M, Dellas C, Lankeit M, Olschewski M, Binder L, Geibel A et al. Heart-type fatty acid-binding protein permits early risk stratification of pulmonary embolism. *Eur Heart J* 2007;**28**:224–229.
- Kostrubiec M, Pruszczyk P, Bochowicz A, Pacho R, Szulc M, Kaczynska A et al. Biomarker-based risk assessment model in acute pulmonary embolism. *Eur Heart J* 2005;**26**:2166–2172.
- Kucher N, Wallmann D, Carone A, Windecker S, Meier B, Hess OM. Incremental prognostic value of troponin I and echocardiography in patients with acute pulmonary embolism. *Eur Heart J* 2003;**24**:1651–1656.
- Scridon T, Scridon C, Skali H, Alvarez A, Goldhaber SZ, Solomon SD. Prognostic significance of troponin elevation and right ventricular enlargement in acute pulmonary embolism. *Am J Cardiol* 2005;**96**:303–305.