

Table S21 Comparison of CBM-containing protein families in *P. tritici-repentis* and other grass and non-grass plant pathogens

Species	CBM family															Total	
	1	6	12	13	18	20	21	24	32	35	38	42	43	48	50		52
<i>Magnaporthe grisea</i> 70-15	22	2	0	0	33	3	1	0	0	3	0	1	1	1	21	1	89
<i>Pyrenophora tritici-repentis</i> Pt-1C-BFP	11	1	0	0	19	2	1	0	1	3	0	1	2	3	0	0	44
<i>Phaeosphaeria nodorum</i> SN15 *	13	1	0	0	43	3	1	0	1	3	0	1	2	2	0	0	70
<i>Gibberella zeae</i> PH-1 (<i>F. graminearum</i>)	12	1	0	2	35	2	2	1	4	2	0	1	1	1	33	0	97
<i>Gibberella moniliformis</i> 7600 (<i>F. verticillioides</i>)	14	1	0	1	26	2	1	1	3	4	1	1	1	3	52	0	111
<i>Fusarium oxysporum</i> f. sp. <i>lycopersici</i> 4286	13	1	4	1	36	2	10	2	3	4	3	1	0	1	46	0	127
<i>Verticillium albo-atrum</i> VaMs.102	26	0	0	0	29	4	1	3	0	3	1	2	1	1	33	0	104
<i>Verticillium dahliae</i> VdLs.17	30	0	0	0	29	4	0	3	0	3	1	2	1	1	32	0	106

* Data from Anselem et al. 2011