

Genotyping of Endosperms to Determine Seed Dormancy Genes Regulating Germination through Embryonic, Endospermic or Maternal Tissues in Rice

Xing-You Gu¹, Jinfeng Zhang², Heng Ye, Lihua Zhang, and Jiuhuan Feng

Plant Science Department, South Dakota State University, Brookings, SD 57007

²Present address: Plant Gene Engineering Lab, DBN Agr. Sci&Tech Research Institute, Beijing 100085, China

¹Corresponding author: Xing-You Gu; mailing address: Box 2140C, Brookings, SD 57007, USA; phone: +1(605) 688-6908; fax: +1(605) 688-4452; email: Xingyou.gu@sdstate.edu

DOI: 10.1534/g3.114.015362

Table S1 Analysis of variance for the time period of incubation (d) required for individual seeds to germinate in the germinated subpopulation segregating for the SD_{7-1} and SD_{12} loci (refer to “ SD_{7-1} & SD_{12} Ex. #1” in Table 3 for additional information)

Source of variance	D.F.	SS	MS	F-value	Probability
SD_{12}	2	88.28	44.141	10.61	<0.0001
SD_{7-1}	2	3.77	1.887	0.45	0.63
$SD_{12} \times SD_{7-1}$	4	18.63	4.657	1.12	0.34
Error	427	1777.1	4.162		
Total	435	1910			