

Literature Cited

- Babu, R., B. M. Prasanna, 2014, Molecular breeding for quality protein maize (QPM) pp 489–505 in *Genomics of plant genetic resources*, edited by R. Tuberosa, A. Graner, E. Frison. Springer, Heidelberg.
- Bailey-Serres, J., T. Fukao, P. Ronald, A. Ismail, S. Heuer et al., 2010 Submergence tolerant rice: SUB1's journey from landrace to modern cultivar. *Rice* 3: 138–147.
- Beckert, P., M. Brinkmann, M. Beckert, 2008 A major locus expressed in the male gametophyte with incomplete penetrance is responsible for in situ gynogenesis in maize. *Theor. Appl. Genet.* 117: 581–594.
- Borojevic, K., and K. Borojevic, 2005 The transfer and history of “reduced height genes” (Rht) in wheat from Japan to Europe. *Journal of Heredity* 96: 455–459.
- Chalyk, S. T., 1999 Creating new haploid-inducing lines of maize. *Maize Genet Coop Newsletter* 73: 53–54.
- Chase, S. S., 1949 Monoploid frequencies in a commercial double cross hybrid maize, and its component single cross hybrids and inbred lines. *Genetics* 34: 328–332.
- Chase, S. S., 1969 Monoploids and monoploid-derivatives of maize (*Zea mays L.*). *Bot. Rev.* 35: 117–167.
- Chia, J.-M., C. Song, P. J. Bradbury, D. Costich, N. de Leon et al., 2012 Maize HapMap2 identifies extant variation from a genome in flux. *Nature Genetics* 44: 803–807.
- Chumak, M. V., 1979 The use of haploid in breeding maize. Proceedings of the tenth meeting of the maize and sorghum section of EUCARPIA. Varna - 17 - 19 - September.
- Clopper, C., E.S. Pearson, 1934 The use of confidence or fiducial limits illustrated in the case of the binomial. *Biometrika* 26: 404–413.
- Coe, E.H., 1959 A line of maize with high haploid frequency. *Am. Nat.* 93: 381–382.
- Consortium, T. U., 2014 UniProt: a hub for protein information. *Nucleic Acids Research* 43: D204–D212.
- Cook, D. E., T. G. Lee, X. Guo, S. Melito, K. Wang et al., 2012 Copy number variation of multiple genes at Rhg1 mediates nematode resistance in soybean. *Science (New York, N.Y.)* 338: 1206–9.
- Cook, J. P., M. D. McMullen, J. B. Holland, F. Tian, P. Bradbury et al., 2012 Genetic Architecture of Maize Kernel Composition in the Nested Association Mapping and Inbred Association Panels. *Plant Physiology* 158: 824–834.
- Delourme, R., C. Falentin, B. F. Fomeju, M. Boillot, G. Lassalle et al., 2013 High-density SNP-based genetic map development and linkage disequilibrium assessment in *Brassica napus L.* *BMC genomics* 14: 120.
- Eder, J., and S. Chalyk, 2002 In vivo haploid induction in maize. *Theoretical and Applied Genetics* 104: 703–708.
- Flint-Garcia, S. A., A. C. Thuitillet, J. Yu, G. Pressoir, S. M. Romero et al., 2005 Maize association population: A high-resolution platform for quantitative trait locus dissection. *Plant Journal* 44: 1054–1064.
- Ganal, M. W., G. Durstewitz, A. Polley, A. Bérard, E. S. Buckler et al., 2011 A large maize (*zea mays L.*) SNP genotyping array: Development and germplasm genotyping, and genetic mapping to compare with the B73 reference genome. *PLoS ONE* 6:.

- Geiger, H. H., and G. A. Gordillo, 2009 Doubled haploids in hybrid maize breeding. *Maydica* 54: 485–499.
- Gerdes, J. T., C. F. Behr, J. G. Coors, W. F. Tracy, 1993 *Compilation of North American Maize Breeding Germplasm*. Crop Sci Soc of Am. Madison, USA.
- Hanson, B. D., L. Fandrich, D. L. Shaner, P. Westra, and S. J. Nissen, 2007 Recovery of imidazolinone-resistant hard red wheat lines following imazamox application. *Crop Science* 47: 2058–2066.
- Jiao, Y., H. Zhao, L. Ren, W. Song, B. Zeng et al., 2012 Genome-wide genetic changes during modern breeding of maize. *Nature Genetics* 44: 812–815.
- Lashermes, P., M. Beckert, 1988 Genetic control of maternal haploidy in maize (*Zea mays* L.) and selection of haploid inducing lines. *Theor Appl Genet* 76: 405–410.
- Liu, Z., T. Song, 2000 The breeding and identification of haploid inducer with high frequency parthenogenesis in Maize. *ACTA AGRONOMICA SINICA* 26: 570–574.
- MBS Inc, 2000 *Genetics Handbook*. Ames, Iowa, USA.
- Melchinger, A. E., W. Schipprack, T. Würschum, S. Chen, and F. Technow, 2013 Rapid and accurate identification of in vivo-induced haploid seeds based on oil content in maize. *Scientific reports*, Nature 3: 1–5.
- Mitchell, a., H.-Y. Chang, L. Daugherty, M. Fraser, S. Hunter et al., 2014 The InterPro protein families database: the classification resource after 15 years. *Nucleic Acids Research* 43: D213–D221.
- Nanda, D. K., and S. S. Chase, 1966 An Embryo Marker for Detecting Monoploids Of Maize (*Zea Mays* L.). *Crop Science* 6: 213.
- Nelson, P. T., 2009 Genetic and phenotypic characterization of maize germplasm resources: Ex-PVPA inbreds, NCSU inbreds, and elite exotic inbreds. PhD dissertation of North Carolina State University, Raleigh, North Carolina.
- Prigge, V., W. Schipprack, G. Mahuku, G. N. Atlin, and A. E. Melchinger, 2012a Development of in vivo haploid inducers for tropical maize breeding programs. *Euphytica* 185: 481–490.
- Prigge, V., X. Xu, L. Li, R. Babu, S. Chen et al., 2012b New Insights into the Genetics of in Vivo Induction of Maternal Haploids , the Backbone of Doubled. *Genetics* 190: 781–793.
- Randolph, L. F., 1940 Note on haploid frequencies. *Maize Genet Coop Newsletter* 14: 23–24.
- Röber, F. K., G. A. Gordillo, and H. H. Geiger, 2005 In vivo haploid induction in maize - Performance of new inducers and significance of doubled haploid lines in hybrid breeding. *Maydica* 50: 275–283.
- Romay, M. C., M. J. Millard, J. C. Glaubitz, J. A. Peiffer, K. L. Swarts et al., 2013 Comprehensive genotyping of the USA national maize inbred seed bank. *Genome biology* 14: R55.
- Rotareenco, V., G. Dicu, D. State, S. Fuia, 2010 New inducers of maternal haploids in maize. *Maize Genet Coop Newsletter* 84: 21–22.
- Sarkar, K. R., A. Pandey, P. Gayen, J.K. Madan, R. Kumar et al., 1994 Stabilization of high haploid inducer lines. *Maize Genet Coop Newsletter* 68: 64–65.
- Septiningsih, E. M., A. M. Pamplona, D. L. Sanchez, C. N. Neeraja, G. V. Vergara et al., 2009 Development of submergence-tolerant rice cultivars: The Sub1 locus and beyond. *Annals of Botany* 103: 151–160.

- Shatskaya, O. A., 2010 Haploinductors isolation in maize: three cycles of selection on high frequency of induction of matroclinal haploids. Agricultural Biology 5: 79–86.
- Snowdon, R. J., W. Lühs, W .Friedt, 2006 Oilseed rap, pp 55–114 in *Genome Mapping and Molecular Breeding in plants*, edited by C. Kole. Springer, Heidelberg.
- Strigens, A., C. Grieder, B. I. G. Haussmann, and A. E. Melchinger, 2012 Genetic variation among inbred lines and testcrosses of maize for early growth parameters and their relationship to final dry matter yield. Crop Science 52: 1084–1092.
- Tyrnov, V. S., A. N. Zavalishina, 1984 Induction of high frequency occurrence of matroclinal haploids in maize. Report of USSR Academy of Science. Genetics 276: 735–738.
- Xu, X., L. Li, X. Dong, W. Jin, A. E. Melchinger et al., 2013 Gametophytic and zygotic selection leads to segregation distortion through *in vivo* induction of a maternal haploid in maize. J. Exp. Botany 64: 1083–1096.
- Xu, K., X. Xu, T. Fukao, P. Canlas, R. Maghirang-Rodriguez et al., 2006 Sub1A is an ethylene-response-factor-like gene that confers submergence tolerance to rice. Nature 442: 705–708.
- Yang, X., S. Gao, S. Xu, Z. Zhang, B. M. Prasanna et al., 2011 Characterization of a global germplasm collection and its potential utilization for analysis of complex quantitative traits in maize. Molecular Breeding 28: 511–526.
- Zabirova, J.R., M.V. Chumak, O. A. Shatskaya, V. S. Scherbak, 1996 Technology of rapid mass generation of homozygous lines. Kukuruza i sorgo 4: 17–19.
- Zavalishina, A. N., V. S. Tyrnov, 1992 Induction of matroclinal haploidy in maize *in vivo*, pp221–222 in XIIIth Eucarpia Congress, Reproductive Biology and Plant Breeding. Angers, France.