Supplementary Material

to

Conservation of anatomically restricted glycosaminoglycan structures in divergent nematode species

Matthew Attreed¹, Kristian Saied-Santiago¹ and Hannes E. Bülow^{1,2}

¹Department of Genetics and ^{1,2}Dominick P. Purpura Department of Neuroscience

Albert Einstein College of Medicine

Bronx, New York, 10461

* corresponding author:

Telephone 718 430 3621

Fax 718 430 8778

e-mail: <u>hannes.buelow@einstein.yu.edu</u>

Running title: conserved glycosaminoglycan structures

3 Figures

1 Table

2 Supplementary Figures

2 Supplementary Tables

Supplementary Materials

<u>Mutants:</u> hse-5(tm472) III (OH1487), hst-2(ok595) X (OH1876), hst-6(ok273) X (OH1421), hst-3.1(tm734) II (EB649), hst-3.2(dz171) X.

Integrated transgenic strains:

EB1119: dzls12 [Punc-122::HS3A8::GFP(S65T); pRF4] EB669: dzls15 [Punc-122::EW4E9::GFP(S65T); pRF4] EB703: dzls16 [Punc-122::MPB49::GFP(S65T); pRF4] (EB703) EB1968: dzEx1031 [Punc-122::EW3G6::2xsfGFP; pRF4] EB1981: dzEx1042 [Punc-122::EW4A4::2xsfGFP; pRF4] EB1782: dzEx876 [Punc-122::EW4A4::2xGFP; Pttx-3::mCherry; pRF4] EB1350: dzEx597 [Punc-122::EW3G6::2xsfgfp; Pglr-6::mCherry; dpy-20(+)] EB1974: dzEx1037 [Punc-122::EW3G6::2xsfGFP; Pgcy-13::mCherry; pRF4] EB2226: dzEx1249 [Punc-122::EW4A4::2xsfGFP; Pinx-1::mCherry; pRF4] EB2228: dzEx1251 [Punc-122::EW4A4::2xsfGFP; Pgcy-13::mCherry; pRF4].

Extrachromosomal transgenes for surveying 33 different HS-specific scFv antibodies (if not mentioned elsewhere):

EB889-90: dzEx468-9 [Punc-122::EW4E10::GFP(S65T);pRF4] EB1408-10: dzEx625-7 [Punc-122::HS3B7::GFP(S65T);pRF4] EB1406-7: dzEx623-4 [Punc-122::EW4A4::GFP(S65T);pRF4] EB1475: dzEx671 [Punc-122::EW4B7::GFP(S65T);pRF4] EB1404-5: dzEx621-2 [Punc-122::HS3G8::GFP(S65T);pRF4] EB888: dzEx467 [Punc-122::EW4B10::GFP(S65T);pRF4] EB604-5: dzEx363-4 [Punc-122::RB4EA12::GFP(S65T);pRF4] EB550-1: dzEx326-7 [Punc-122::EV3B2::GFP(S65T);pRF4] EB539-42: dzEx315-8 [Punc-122::EW3D3::GFP(S65T);pRF4] EB552-3: dzEx328-9 [Punc-122::EW3E4::GFP(S65T);pRF4] EB554-6: dzEx330-2 [Punc-122::EW3F5::GFP(S65T);pRF4] EB571: dzEx343 [Punc-122::EW3G6::GFP(S65T);pRF4] EB916-8: dzEx474-6 [Punc-122::EW3G6::2xsfGFP;pRF4] EB1459: dzls34 [Punc-122::EW3G6::2xsfGFP;pRF4] EB1969: *dzEx1032* [*Punc-122::EW3G6::2xsfGFP;pRF4*] EB547-9: dzEx323-5 [Punc-122::EW4A11::GFP(S65T);pRF4] EB543-6: dzEx319-22 [Punc-122::EW4E1::GFP(S65T);pRF4] EB532-3: dzEx308-9 [Punc-122::EW4E9::GFP(S65T);pRF4] EB757: dzls15 [Punc-122::EW4E9::GFP(S65T);pRF4] EB1114: dzEx519 [Punc-122::EW4E9::GFP(S65T);pRF4] EB574-6: dzEx344-6 [Punc-122::EW4G2::GFP(S65T);pRF4] EB1204-8: dzls27-31 [Punc-122::EW4G2::GFP(S65T);pRF4] EB1325, EB1327: dzls32-3 [Punc-122::EW4G2::GFP(S65T);pRF4] EB568-70: dzEx340-2 [Punc-122::HS3A8::GFP(S65T);pRF4] EB666-7: dzls12-3 [Punc-122::HS3A8::GFP(S65T);pRF4] EB313-9: dzEx195-201 [Punc-122::HS4C3::GFP(S65T);pRF4] EB502-3: dzls10-1 [Punc-122::HS4C3::GFP(S65T);pRF4] EB536-7: dzEx312-4 [Punc-122::HS4E4::GFP(S65T);pRF4] EB1104: dzEx511 [Punc-122::HS4E4::GFP(S65T);pRF4] EB1105: dzEx512 [Punc-122::LKIV69::GFP(S65T);pRF4] EB1202-3: dzls25-6 [Punc-122::LKIV69::GFP(S65T);pRF4] EB1333-4: dzEx580-1 [Punc-122::EV3C3::GFP(S65T);pRF4] EB1399-403: dzEx616-620 [Punc-122::EW4G1::GFP(S65T);pRF4] EB1110-1: dzEx516-7 [Punc-122::HS4D10::GFP(S65T);pRF4]

EB1472-3: *dzEx668-9* [*Punc-122::EW4B5::GFP*(S65T);*pRF4*] EB1474,1601-4: *dzEx670,760-3* [*Punc-122::EW4C10::GFP*(S65T);*pRF4*] EB749-50,891-2: *dzEx403-4,470-1* [*Punc-122::EW4D5::GFP*(S65T);*pRF4*] EB1476-7: *dzEx672-3* [*Punc-122::EW4G10::GFP*(S65T);*pRF4*] EB636-8: *dzEx368-370* [*Punc-122::AO4B08::GFP*(S65T);*pRF4*] EB668: *dzls14* [*Punc-122::AO4B08::GFP*(S65T);*pRF4*] EB534-5,664-5: *dzEx310-1,376-7* [*Punc-122::EV3D1::GFP*(S65T);*pRF4*] EB1398: *dzEx615* [*Punc-122::MW3G3::GFP*(S65T);*pRF4*] EB1723: *dzEx834* [*Punc-122::NS4F5::GFP*(S65T);*pRF4*] EB1595: *dzEx759* [*Punc-122::HS4D4::GFP*(S65T);*pRF4*] EB1347-9,1724-5: *dzEx594-6,835-6* [*Punc-122::HS4A5::GFP*(S65T);*pRF4*]

Transgenes for cell identification of EW3G6:

EB1106-7: *dzEx513-4 [Punc-122::ew3g6::2xgfp 25ng/µl; Pttx-3::mCherry 75ng/µl; pRF4 75ng/µl].* Similar DNA concentrations were used in all following strains, except for strains with complex arrays (identifiable as containing gDNA N2).

EB1121-2: dzEx518+dzEx520 [Punc-122::ew3g6::2xgfp; Pttx-3::mCherry; pRF4] EB1772-3: dzEx866-7 [Punc-122::EW3G6::2xGFP; Pflp-8::mCherry; pRF4] EB1344: dzEx591 [Punc-122::ew3g6::2xsfgfp; Ppept-3::mCherry; dpy-20(+)] EB1353: dzEx599 [Punc-122::ew3g6::2xsfgfp; Ppept-3::mCherry; dpy-20(+)] EB1338: dzEx585 [Punc-122::ew3g6::2xsfGFP; Pceh-16::mCherry; dpy-20(+)] EB1339: dzEx586 [Punc-122::ew3g6::2xsfGFP; Pceh-16::mCherry; dpy-20(+)] EB1330: otls181; dzEx475 [Punc-122::ew3g6::2xsfGFP] EB1329: otls266; dzEx475 [Punc-122::ew3g6::2xsfGFP] EB1350-1: dzEx597-8 [Punc-122::ew3g6::2xsfgfp; Pglr-6::mCherry; dpy-20(+)] EB1774-5:dzEx868-9 [Punc-122::EW3G6::2xGFP; Ptdc-1::mCherry; pRF4] EB1768-9: dzEx862-3 [Punc-122::EW3G6::2xGFP; Pzig-3::mCherry; pRF4]

EB1770-1: dzEx864-5 [Punc-122::EW3G6::2xGFP; Psra-11::mCherry; pRF4]

EB1972-3: dzEx1035-6 [complex array Punc-122::ew3g6::2xsfGFP; Ptbh-1::mCherry PCR fusion; pRF4; N2 gDNA]

EB1976: dzEx1039 [complex array Punc-122::ew3g6::2xsfGFP; Pgcy-28::mCherry PCR fusion; pRF4; N2 gDNA]

EB1974-5: dzEx1037-8 [complex array Punc-122::ew3g6::2xsfGFP; Pgcy-13::mCherry PCR fusion; pRF4; N2 gDNA]

EB1970-1: dzEx1033-4 [complex array Punc-122::ew3g6::2xsfGFP; Pinx-1::mCherry PCR fusion; pRF4; N2 gDNA]

EB2299-2300: dzEx1285-6 [complex array: Punc-122::ew3g6::gfp Pgcy-18::mCherry; pRF4; N2 gDNA]

Transgenes for cell identification of EW4A4:

EB1592-4: dzls34; dzEx756-8 [Punc-122::ew4a4::DsRed2; Pttx-3::mCherry; pBS] EB1782-3: dzEx876-7 [Punc-122::EW4A4::2xGFP; Pttx-3::mCherry; pRF4] EB1776-7:dzEx870-1 [Punc-122::EW4A4::2xGFP; Pser-2.2::mCherry; pRF4] EB1778: dzEx872 [Punc-122::EW4A4::2xGFP; Pflp-8::mCherry; pRF4] EB1779: dzEx873 [Punc-122::EW4A4::2xGFP; Pfax-1::mCherry; pRF4] EB1784-5:dzEx878-9 [Punc-122::EW4A4::2xGFP; Ptdc-1::mCherry; pRF4] EB1780-1:dzEx874-5 [Punc-122::EW4A4::2xGFP; Pgcy-32::mCherry; pRF4] EB2228-9:dzEx1251-2 [complex array: Punc-122::EW4A4::2xsfGFP; Pgcy-13::mCherry; pRF4; N2 gDNA]

EB2226-7: dzEx1249-50 [complex array: Punc-122::EW4A4::2xsfGFP; Pinx-1::mCherry; pRF4; N2 gDNA]

Transgenes in the C. elegans CB4856 strain:

EB1613: *dzEx772* [*Punc-122::EW4A4::2xsfGFP; pRF4*] EB1618: *dzEx777* [*Punc-122::EW3G6::2xsfGFP; pRF4*]

<u>Transgenes in C. briggsae (AF16):</u> dzEx1024 [Punc-122::HS3A8::GFP(S65T)] dzEx1025 [Punc-122::EW4A4::2xsfGFP; pRF4] dzEx1026 [Punc-122::EW3G6::2xsfGFP; pRF4].

Supplementary Figures

Figure S1. Triangulation with cell specific markers was used to identify the location of the specific label in EW4A4::2xsfGFP and EW3G6::2xsfGFP transgenic animals.

- a. c'. Epifluorescent optical sections of a medial/sagittal (a-c) and a lateral/parasagittal plane (a'-c') of animals transgenically expressing the EW4A4::2xsfGFP scFv antibody fusion and mCherry as a cytoplasmic marker in the bilateral pair of AIYL/R interneurons under control of the *Pttx-3* promoter (Altun-Gultekin, Z., Andachi, Y., et al. 2001). Images of the green (a,a') and red (b,b') channels with a merged image (c,c') are shown. Green and red arrowheads point to labeling of the respective processes. Anterior is to the left, ventral down in all images and a scale bar indicates 10 µm in all images (a-f').
- b. f'. Epifluorescent optical sections of a medial/sagittal (d-f) and a lateral/parasagittal plane (d'-f') of animals transgenically expressing the EW3G6::2xsfGFP scFv antibody fusion and mCherry as a cytoplasmic marker in the bilateral pair of RIAL/R interneurons under control of the *Pglr-6* promoter (Brockie, P.J., Madsen, D.M., et al. 2001). Images of the green (d,d') and red (e,e') channels with a merged image (f,f') are shown. Green and red arrowheads point to labeling of the respective processes.
- g. Schematic of the dorsal portion of the *C. elegans* nerve cords from transverse sections with the names of processes indicated. Orange lines exemplify the triangulation strategy using AIY and RIA neurons (indicated in magenta) pointing to the area in the vicinity of AIB and RIM neurons (indicated in green). Modified after Fig.20c in (White, J.G., Southgate, E., et al. 1986).

Figure S2. EW3G6::2xsfGFP and EW4A4::2xsfGFP bind distinct epitopes in proximity to the synaptic connection between AIB and RIM interneurons in the *C. elegans* nerve ring.

- a. b. Schematics of RIM interneurons (left panel) and AIB motor/interneurons (right panel) with synaptic output and input between both, respectively, indicated by blue arrows. Modified after <u>www.wormatlas.org</u>.
- c. e'. Epifluorescent optical sections of a medial/sagittal (c,d,e) and a lateral/parasagittal plane (c',d',e') of animals transgenically expressing the EW3G6::2xsfGFP and mCherry as a cytoplasmic marker in RIML/R under control of the *Pgcy-13* promoter (Ortiz, C.O., Etchberger, J.F., et al. 2006). Similar results were obtained with the *Ptdc-1* promoter, which is also expressed in RIM (Alkema, M.J., Hunter-Ensor, M., et al. 2005). Images of the green (c,c') and red (d,d') channels with a merged image (e,e') are shown. Green and red arrowheads point to labeling of the respective processes. Anterior is to the left, ventral down and a scale bar indicates 10 µm in all panels (c-h'). The reciprocal experiment between EW4A4 and AIBL/R is shown in Fig.2.
- f. h'. Epifluorescent optical sections of a medial/sagittal (f,g,h) and a lateral/parasagittal plane (f',g',h') of animals transgenically expressing the EW4A4::2xsfGFP and mCherry as a cytoplasmic marker in the bilateral pair AIBL/R under control of the *Pinx-1* promoter (Altun, Z.F., Chen, B., et al. 2009). Images of the green (f,f') and red (g,g') channels with a merged image (h,h') are shown. Green and red arrowheads point to labeling of the respective processes and indicate the more anterior descending (a), and posterior ascending (p) neurites of AIB, respectively. The reciprocal experiment between EW3G6 and RIML/R is shown in Fig.2.

 Schematic representations of the synaptic connectivity of AIBL and RIMR, modified from (White, J.G., Southgate, E., et al. 1986). Synaptic output to and input from, are indicated by blue arrows for AIBL and RIMR, respectively.

References

Alkema MJ, Hunter-Ensor M, Ringstad N, Horvitz HR. 2005. Tyramine Functions independently of octopamine in the Caenorhabditis elegans nervous system. *Neuron*, 46:247-260.

Altun ZF, Chen B, Wang ZW, Hall DH. 2009. High resolution map of Caenorhabditis elegans gap junction proteins. *Developmental dynamics : an official publication of the American Association of Anatomists*, 238:1936-1950.

Altun-Gultekin Z, Andachi Y, Tsalik EL, Pilgrim D, Kohara Y, Hobert O. 2001. A regulatory cascade of three homeobox genes, ceh-10, ttx-3 and ceh-23, controls cell fate specification of a defined interneuron class in C. elegans. *Development*, 128:1951-1969.

Attreed M, Desbois M, van Kuppevelt TH, Bülow HE. 2012. Direct visualization of specifically modified extracellular glycans in living animals. *Nature methods*, 9:477-479.

Aurelio O, Hall DH, Hobert O. 2002. Immunoglobulin-domain proteins required for maintenance of ventral nerve cord organization. *Science*, 295:686-690.

Bertrand V, Hobert O. 2009. Linking asymmetric cell division to the terminal differentiation program of postmitotic neurons in C. elegans. *Developmental cell*, 16:563-575.

Brockie PJ, Madsen DM, Zheng Y, Mellem J, Maricq AV. 2001. Differential expression of glutamate receptor subunits in the nervous system of Caenorhabditis elegans and their regulation by the homeodomain protein UNC-42. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 21:1510-1522.

Dennissen MA, Jenniskens GJ, Pieffers M, Versteeg EM, Petitou M, Veerkamp JH, van Kuppevelt TH. 2002. Large, tissue-regulated domain diversity of heparan sulfates demonstrated by phage display antibodies. *The Journal of biological chemistry*, 277:10982-10986.

Didiano D, Cochella L, Tursun B, Hobert O. 2010. Neuron-type specific regulation of a 3'UTR through redundant and combinatorially acting cis-regulatory elements. *Rna*, 16:349-363.

Fei YJ, Romero MF, Krause M, Liu JC, Huang W, Ganapathy V, Leibach FH. 2000. A novel H(+)-coupled oligopeptide transporter (OPT3) from Caenorhabditis elegans with a predominant function as a H(+) channel and an exclusive expression in neurons. *The Journal of biological chemistry*, 275:9563-9571.

Hobert O. 2002. PCR fusion-based approach to create reporter gene constructs for expression analysis in transgenic C. elegans. *BioTechniques*, 32:728-730.

Inada H, Ito H, Satterlee J, Sengupta P, Matsumoto K, Mori I. 2006. Identification of guanylyl cyclases that function in thermosensory neurons of Caenorhabditis elegans. *Genetics*, 172:2239-2252.

Kim K, Li C. 2004. Expression and regulation of an FMRFamide-related neuropeptide gene family in Caenorhabditis elegans. *J Comp Neurol*, 475:540-550.

Much JW, Slade DJ, Klampert K, Garriga G, Wightman B. 2000. The fax-1 nuclear hormone receptor regulates axon pathfinding and neurotransmitter expression. *Development*, 127:703-712.

Ortiz CO, Etchberger JF, Posy SL, Frokjaer-Jensen C, Lockery S, Honig B, Hobert O. 2006. Searching for neuronal left/right asymmetry: genomewide analysis of nematode receptor-type guanylyl cyclases. *Genetics*, 173:131-149.

Shinkai Y, Yamamoto Y, Fujiwara M, Tabata T, Murayama T, Hirotsu T, Ikeda DD, Tsunozaki M, Iino Y, Bargmann Cl, *et al.* 2011. Behavioral choice between conflicting alternatives is regulated by a receptor guanylyl cyclase, GCY-28, and a receptor tyrosine kinase, SCD-2, in AIA interneurons of Caenorhabditis elegans. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 31:3007-3015.

Smits NC, Kurup S, Rops AL, ten Dam GB, Massuger LF, Hafmans T, Turnbull JE, Spillmann D, Li JP, Kennel SJ, *et al.* 2010. The heparan sulfate motif (GlcNS6S-IdoA2S)3, common in heparin, has a strict topography and is involved in cell behavior and disease. *The Journal of biological chemistry*, 285:41143-41151.

Smits NC, Lensen JF, Wijnhoven TJ, Ten Dam GB, Jenniskens GJ, van Kuppevelt TH. 2006. Phage display-derived human antibodies against specific glycosaminoglycan epitopes. *Methods Enzymol*, 416:61-87.

ten Dam GB, van de Westerlo EM, Smetsers TF, Willemse M, van Muijen GN, Merry CL, Gallagher JT, Kim YS, van Kuppevelt TH. 2004. Detection of 2-O-sulfated iduronate and N-acetylglucosamine units in heparan sulfate by an antibody selected against acharan sulfate (IdoA2S-GlcNAc)n. *The Journal of biological chemistry*, 279:38346-38352.

Tsalik EL, Niacaris T, Wenick AS, Pau K, Avery L, Hobert O. 2003. LIM homeobox genedependent expression of biogenic amine receptors in restricted regions of the C. elegans nervous system. *Developmental biology*, 263:81-102.

van de Westerlo EM, Smetsers TF, Dennissen MA, Linhardt RJ, Veerkamp JH, van Muijen GN, van Kuppevelt TH. 2002. Human single chain antibodies against heparin: selection, characterization, and effect on coagulation. *Blood*, 99:2427-2433.

van Kuppevelt TH, Dennissen MA, van Venrooij WJ, Hoet RM, Veerkamp JH. 1998. Generation and application of type-specific anti-heparan sulfate antibodies using phage display technology. Further evidence for heparan sulfate heterogeneity in the kidney. *The Journal of biological chemistry*, 273:12960-12966.

White JG, Southgate E, Thomson JN, Brenner S. 1986. The structure of the nervous system of the nematode *Caenorhabditis elegans*. *Philosophical Transactions of the Royal Society of London B. Biological Sciences*, 314:1-340.

Wijnhoven TJ, van de Westerlo EM, Smits NC, Lensen JF, Rops AL, van der Vlag J, Berden JH, van den Heuvel LP, van Kuppevelt TH. 2008. Characterization of anticoagulant heparinoids by immunoprofiling. *Glycoconjugate journal*, 25:177-185.

Yu S, Avery L, Baude E, Garbers DL. 1997. Guanylyl cyclase expression in specific sensory neurons: a new family of chemosensory receptors. *Proceedings of the National Academy of Sciences of the United States of America*, 94:3384-3387.

		•		Н	HS modifications required for binding ^a							а	anatomical binding in <i>C. elegans</i> ^b									
						in v	itro)		In vivo					nervous system				alime sys	ntary tem		
Name of scFv	V _H c	CDR3	isolat. against [₫]	NAc	NS SN	C5 26	22	6S 3<	NAC	NS	C5	2S	6S	3S	nr	vnc	dnc	con	nments	pm, pi valve	gut	refs. ^e
EV3B2 (EW4D2) EW3D3	DP-38 DP-38	GKMKLNR GRTVGRN	hIHS/ iHep iHep		+	+ +	÷	++							++	++	+					(Dennissen, M.A., Jenniskens, G.J., et al. 2002, van de Westerlo, E.M., Smetsers, T.F., et al. 2002) (van de Westerlo, E.M., Smetsers, T.F., et al. 2002)
(EW3D10) EW4E1	DP-38	GRRHKLIR	iHep												+	+	+					(van de Westerlo, E.M., Smetsers, T.F., et al. 2002)
EW4B5	DP-45	GRLHLPRK	іНер												+	+	+					(van de Westerlo, E.M., Smetsers, T.F., et al. 2002)
EW4D5	DP-45	PVSHRKWRVTV	iHep												+							(van de Westerlo, E.M., Smetsers, T.F., et al. 2002)
HS4C3	DP-38	GRRLKD	bkHS		+	(+	+)	+ +				(+)	+	+	+	+	+				(+)	(Attreed, M., Desbois, M., et al. 2012, Smits, N.C., Lensen, J.F.,
HS3A8 (EW3H12)	DP-38	GMRPRL	bkHS		+	+ +	+	+				+	+		+	+	+				(+)	(Attreed, M., Desbois, M., et al. 2012, Dennissen, M.A., Jenniskens, G.J., et al. 2002, van de Westerlo, E.M., Smetsers, T.F., et al. 2002, Wijnhoven, T.J., van de Westerlo, E.M., et al. 2008)
EW4G2	DP-38	GKVKLPN	іНер		+			+							+	+	+					(van de Westerlo, E.M., Smetsers, T.F., et al. 2002)
EW3F5	DP-38	SGRQARQGRFPK	іНер												+	+	+					(van de Westerlo, E.M., Smetsers, T.F., et al. 2002)
HS4E4	DP-38	HAPLRNTRTNT	bkHS	+	+	+ +	+	in.							+	+	+					(Dennissen, M.A., Jenniskens, G.J., et al. 2002, Wijnhoven, T.J., van de Westerlo, E.M., et al. 2008)
EW4C10	DP-45	ARMTGHVRNVMI	іНер												+	+	+					(van de Westerlo, E.M., Smetsers, T.F., et al. 2002)
EW4G1	DP-42	GARLKR	іНер												+	+	+				(+)	(van de Westerlo, E.M., Smetsers, T.F., et al. 2002)
AO4B08 ^r	DP-47	SLRMNGWRAHQ	mskHS		+	+ +	ł	+				+	+		+	+	+			+		(Attreed, M., Desbois, M., et al. 2012, Dennissen, M.A., Jenniskens, G.J., et al. 2002, Wijnhoven, T.J., van de Westerlo, F.M. et al. 2008)
RB4EA12	DP-32	RRYALDY	hsmHS	+	+	+ ir	٦.	+							+							(Dennissen, M.A., Jenniskens, G.J., et al. 2002, Wijnhoven, T.J., van de Westerlo, E.M., et al. 2008)
LKIV69	DP-38	GSRSSR	bkHS		+	+ +	+								+	(+)	(+))		+		(Wijnhoven, T.J., van de Westerlo, E.M., et al. 2008)
HS4D4	DP-58	GMRPRL	bkHS		+	+ +	+	+ (+	·)						+							(Dennissen, M.A., Jenniskens, G.J., et al. 2002)
EW4A4	DP-08	GTKLKMTK	іНер	-								+	+	+ ^g	+			sing	gle neuron			(van de Westerlo, E.M., Smetsers, T.F., et al. 2002), this study.
EW3G6	DP-38	GGTTRIRK	іНер	(+)								+	in.	+/- ^r	+			sing	gle neuron			(van de Westerlo, E.M., Smetsers, T.F., et al. 2002), this study.
EW4E9	DP-38	LRGTKMFRH	іНер															Var	. staining			(van de Westerlo, E.M., Smetsers, T.F., et al. 2002)
EV3C3	DP-42	GYRPRF	hIHS															wea	ak/absent			(Dennissen, M.A., Jenniskens, G.J., et al. 2002)
EV3D1	DP-47	SISMNGVGVRIQ	hIHS	+	+	(+	+)	(+)										wea	ak/absent			(Dennissen, M.A., Jenniskens, G.J., et al. 2002)
EW4E10	DP-03	SRKTPKPFMRK	іНер															wea	ak/absent			(van de Westerlo, E.M., Smetsers, T.F., et al. 2002)
HS3B7	DP-03	SRKTRKPFMRK	bkHS	+	+	+ ir	٦.	+										wea	ak/absent			(Dennissen, M.A., Jenniskens, G.J., et al. 2002)
EW4B7	DP-08	SSSRHHRLHR	іНер															wea	ak/absent			(van de Westerlo, E.M., Smetsers, T.F., et al. 2002)
HS3G8	DP-08	YYHYKVN	bkHS ⁱ												1			wea	ak/absent			(van Kuppevelt, T.H., Dennissen, M.A., et al. 1998)
EW4B10	DP-25	QRWKPAVTPKLV	іНер															wea	ak/absent			(van de Westerlo, E.M., Smetsers, T.F., et al. 2002)

Table SI. HS scFv properties and binding characteristics in vitro and in vivo

EW3E4	DP-38 DRRNTQKTRYRT	іНер					weak/absent	(van de Westerlo, E.M., Smetsers, T.F., et al. 2002)
EW4A11	DP-38 ERNTIRR	іНер					weak/absent	(van de Westerlo, E.M., Smetsers, T.F., et al. 2002)
HS4D10	DP-42 SLRMNGCGAHQ	bkHS					weak/absent	(van Kuppevelt, T.H., Dennissen, M.A., et al. 1998)
EW4G10	DP-45 GTKKLGK	іНер					weak/absent	(van de Westerlo, E.M., Smetsers, T.F., et al. 2002)
MW3G3	DP-47 QKKRPRF	AS	+	+	+		weak/absent	(ten Dam, G.B., van de Westerlo, E.M., et al. 2004)
NS4F5	DP-53 SGRKGRMR	hIHS	+	+	+		weak/absent	(Smits, N.C., Kurup, S., et al. 2010)
HS4A5	DP-65 WVTEP	bkHS	+	+	+	+	weak/absent	(Dennissen, M.A., Jenniskens, G.J., et al. 2002)
MPB49	DB-38 WRNDRQ	na					neg. contr.	(van Kuppevelt, T.H., Dennissen, M.A., et al. 1998)

^a HS modifications required for binding of the scFvs based on competitive binding assays (*in vitro*) or as determined genetically (*in vivo*). +, required; (+) weakly required; -, not required, in., inhibitory. ^b Abbreviations used: nr, nerve ring; vnc, ventral nerve cord; dnc, dorsal nerve cord; pi, pharyngeal intestinal; pm, pharyngeal muscle; +, expressed; (+), weakly expressed. ^cV_H: variable heavy chain type, with the sequence of the complementary determining region 3 indicated in the adjacent column CDR3. ^d Target material to isolate scFvs in panning experiments is indicated: iHep, immobilized heparin; bkHS, bovine kidney HS; hsmHS, human skeletal muscle HS; hlHS, human lung HS; mskHS, mouse skeletal muscle HS; AS, acharan sulfate. ^e References are indicated that initially describe isolation, biochemical characterization, or in the case of HS4C3, HS3A8, and AO4B08 prior genetic characterization in *C. elegans*. ^f Note that AO4B08 shows some additional staining in the M-lines of the muscle and the dense bodies. ^g Both type I and II HS 3-O-sulfotransferases are required for binding *in vivo*. ^h HS 3-O-sulfation introduced by type I HS 3-O-sulfotransferases is required for binding *in vivo*. ^h HS 3-O-sulfotransferases is required for binding *in vivo*. ^h HS 3-O-sulfotransferases are required for binding *in vivo*. ^h HS 3-O-sulfotransferases are required for binding *in vivo*. ^h HS 3-O-sulfotransferases are required for binding *in vivo*. ^h HS 3-O-sulfotransferases is required for binding *in vivo*. ^h HS 3-O-sulfotransferases are required for binding *in vivo*. ^h HS 3-O-sulfotransferases is required for binding in *C. elegans* (this study). ⁱ HS3G8 was counterselected against chondroitin and dermatan sulfate.

		tested with	ו				
Geneª	neurons labeled ^ь	EW3G6°	EW4A4 ^c	5'primer used for PCR fusion	3' fusion used for PCR fusion	size	ref ^d
ttx-3*	AIY	yes (4)	yes (5)	na	na		(Altun-Gultekin, Z., Andachi, Y., et al. 2001)
ser-2.2*	AIY AIZ RID DVA BDU SIAD SIAV RME PVT	No	yes (2)	na	na		(Tsalik, E.L., Niacaris, T., et al. 2003)
flp-8	ADA CP9 URX RMG AUA PVM	yes (2)	yes (1)	TCTTCGTGATGGAGTTCGC	ATCCTCCTCGCCCTTGCTCACCATT TTCTACTTGAAAAGTGTGG	3.0kb	(Kim, K. and Li, C. 2004)
opt-3 (pept-3)	AVE	yes (2)	no	CGATAAAAATAACAGAATTAGT AAGAAGGTGGG	ATCCTCCTCGCCCTTGCTCACCATG GTTAATTGAGATTCCTTTTGC	2.8kb	(Fei, Y.J., Romero, M.F., et al. 2000)
ceh-16	GLRD	yes (2)	no	AAGGCGCATAAAATGTATTCG	ATCCTCCTCGCCCTTGCTCACCATA ATACCGAATTTCAGAATCAT	3.1kb	na
dat-1 (otis181)**	PDE CEPV CEPD ADE	yes (1)	no	na	na		(Bertrand, V. and Hobert, O. 2009)
che-1 (otls232)**	ASE ASH AUA	yes (1)	no	na	na		(Didiano, D., Cochella, L., et al. 2010)
fax-1*	AVK AVE SABD AVB CEPD SIBD RIC SMBV SABV AVA AIY URX PVP DVA	no	yes (1)	na	na		(Much, J.W., Slade, D.J., et al. 2000)
glr-6	RIA	yes (2)	no	GGCCCGAGTCACAAATACCGC	ATCCTCCTCGCCCTTGCTCACCATT TCAAATGCATTTGTTCAGC	4.0kb	(Brockie, P.J., Madsen, D.M., et al. 2001)
tdc-1	RIC RIM	yes (2)	yes (2)	AAGTAGTTGGTGTACATAGTTT CGG	ATCCTCCTCGCCCTTGCTCACCATT TGGGCGGTCCTGAAAAATGCACCG	3.2kb	(Alkema, M.J., Hunter-Ensor, M., et al. 2005)
zig-3	PVT AIM ASI	yes (2)	no	GCTCGCCAAGAAACATTCCTTT ACCC	ATCCTCCTCGCCCTTGCTCACCATT TTTCCAAATGCAAAATATAG	4.5kb	(Aurelio, O., Hall, D.H., et al. 2002)
sra-11	AIA AIY AVB RIG RIF	yes (2)	no	ACTAAACTAATTAACAAGTCAC GG	ATCCTCCTCGCCCTTGCTCACCATG GTTAGCTGAAATATAAGGG	4.2kb	(Altun-Gultekin, Z., Andachi, Y., et al. 2001)
gcy-32	URX AQR PQR	no	yes (2)	GTTTTCCAAAGGCGAAACGTGG GG	ATCCTCCTCGCCCTTGCTCACCATT CTATAATACAATCGTGATCTTCGC	4.1kb	(Yu, S., Avery, L., et al. 1997)
tbh-1	RIC	yes (2)	no	GCCGCGTTGGCCGCAATTTT	ATCCTCCTCGCCCTTGCTCACCATA GCAACGGCACTTCTCAT	5.0kb	(Alkema, M.J., Hunter-Ensor, M., et al. 2005)
gcy-28	AIA	yes (1)	no	TACAATTGTAGTGAGCTTCG	ATCCTCCTCGCCCTTGCTCACCATT TCGCACTCATCTCACCATTCC	2.8kb	(Shinkai, Y., Yamamoto, Y., et al. 2011)
gcy-13	RIM	yes (2)	yes (2)	CTCAAAATCCATTGTGTAAGTT C	ATCCTCCTCGCCCTTGCTCACCATC CATCCTACAGAGGAGGC	2.3kb	(Ortiz, C.O., Etchberger, J.F., et al. 2006)
inx-1	AIB AIY	yes (2)	yes (2)	CAGATTCATAACTGTCTATG	ATCCTCCTCGCCCTTGCTCACCATC	0.9kb	(Altun, Z.F., Chen, B., et al. 2009)
gcy-18	AFD	yes (2)	no	GGTCGAGAGCAGCAAATC	ATCCTCCTCGCCCTTGCTCACCATT TTCTGATGCTCCGAC	0.8kb	(Inada, H., Ito, H., et al. 2006)

Table SII. Cell specific reporters used for triangulation experiments.

^a Promoters of genes used to drive cytoplasmic mCherry for triangulation experiments with EW3G6-2xsfGFP and EW4A4-2xsfGFP in the nerve ring using existing constructs (*) or integrated reporter lines (**). All other experiments were conducted using promoter fusions obtained by a PCR fusion technique (Hobert, O. 2002) with the primers indicating the extent of the 5'UTR. ^b Labeled neurons as reported on <u>www.wormbase.org</u>. ^c Number of transgenic lines with comparable results indicated in parentheses. ^d References are indicated that either describe the constructs or were used as a basis to design primers. na: not applicable.



