

Supplementary data:

Expression patterns of *FLAGELLIN SENSING 2* map to bacterial entry sites in plant shoots and roots

Martina Beck¹, Ines Wyrsch², James Strutt¹, Rinukshi Wimalasekera³, Alex Webb³, Thomas Boller² and Silke Robatzek¹

¹The Sainsbury Laboratory, Norwich Research Park, Norwich, NR4 7UH, UK

²Zürich-Basel Plant Science Center; University of Basel; Department of Environmental Sciences; Botany; Basel, Switzerland

³Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge CB2 3EA, UK

Supplement Table:

Table 1: Flg22 up-regulated genes in roots, microarray data

Supplement Figures:

Fig S1: Prediction of FLS2 promoter motifs 1000bp upstream of At5g46330

(A) Visualization of promoter motifs 1000 bp upstream of ATG. (B) Details of predicted promoter motifs 1000 bp upstream of ATG.

Prediction with Athena: a resource for rapid visualization and systematic analysis of Arabidopsis promoter sequences (O'Connor *et al.*, 2005; <http://www.bioinformatics2.wsu.edu/Athena>).

Figure S2: FLS2 promoter activity during plant development

pFLS2::GUS is ubiquitously expressed in (A) 2 days-old seedling, in (B) 8 days-old seedling cotyledons and (C) hypocotyl; and highly expressed in (D) flower sepals and (E) stamen, in (F) pod dehiscence zone in mature siliques and (G) stipules.

Figure S3: FLS2 promoter activity during leaf development, wound stress and biotic stress


(A) Definition of different leaf stages; (B) GUS staining of *pFLS2::GUS* in different leaf stages. Insert shows hydathodes. (C) Wound-induced promoter activity in different leaf stages. Insert shows hydathodes. (D) Hypocotyl after *Pto DC3000* incubation with enhanced promoter activity in stomata (arrows), compared to mock treated (10mM MgCl₂) hypocotyl; bar = 100 μm.

Figure S4: *Pto DC3000*-GFP localization on leaves and roots

(A) Confocal micrographs showing *Pto* DC3000-GFP (green) on Col-0 leaves, upper panel shows bacteria on epidermis, encircled are stomata, lower panel shows bacteria (arrow heads) in the apoplast of mesophyll cells (block arrows); chloroplast autofluorescence is represented in red. **(B)** Confocal micrographs showing DC3000-GFP (green) bacteria on Col-0 roots stained with propidium iodide (red); note the accumulation of bacteria (arrow heads) at outgrowing LR (block arrows).

Figure S5: FLS2 promoter activity in non-sterile grown roots and flg22 dependent inhibition of root growth

(A) *pFLS2::GUS* activity in 14 days-old seedlings grown on soil, white arrow shows inner vasculature, black arrows mark expansion of GUS staining, bar = 50 μ m. **(B)** Cross section of roots grown on soil, arrows mark expanded GUS activity in endodermis, bar = 10 μ m. **(C)** Graph depicts root length of 12 days-old Col-0 or *fls2* treated with 1 μ M flg22 or without (same seedlings as used for 3F, bars represent average of 3 independent experiments, error bars represent SD, statistical significance represented with student t-test (p-value >0.001).



B

At5g46330		
Motifs		
AtMYC2 BS in RD22		CACATG
189 -184	(-)	CATGTG
CARGCW8GAT		CWWWWWWWWWG
329 -320	(+)	CATAAATAAG
329 -320	(-)	CATAAATAAG
CCA1 binding site motif		AAMAATCT
826 -819	(-)	AGATTGTT
GAREAT		TAACAAR
593 -587	(+)	TAACAAA
Gap-box Motif		CAAATGAA
962 -955	(-)	TTCATTTG
L1-box promoter motif		TAAATGYA
164 -157	(-)	TACATTAA
LEAFYATAG		CCAATGT
953 -947	(+)	CCAATGT
MYB1AT		WAACCA
710 -705	(+)	AAACCA
MYB4 binding site motif		AMCWAMC
550 -544	(-)	GTTTGT
MYCATERD1		CATGTG
189 -184	(+)	CATGTG
T-box promoter motif		ACTTTG
381 -376	(+)	ACTTG
TATA-box Motif		TATAAA
484 -479	(+)	TATAAA
29 -24	(+)	TATAAA
151 -146	(-)	TTTATA
W-box promoter motif		TTGACY
601 -596	(+)	TTGACC
432 -427	(+)	TTGACT
655 -650	(-)	AGTCAA
EIN3 binding sites		ATGTATCT
325-317		ATGAATaT
304-297		AcATACAT
285-278		ATGAATaT

Figure S1




Figure S2




Figure S3




Figure S4




Figure S5