
Presence	e / absence homology	0
	(mar.)	
	presence c	
	of homologs in organism(	s) :
	Bacillus cereus strain G9842 [taxonId=405531] x	
	Lactococcus lactis strain KF147 [taxonId=684738] x	
	( AND)	9
	absence 🗢	
	of homologs in organsim(	s) :
	Staphylococcus aureus strain ED98 [taxonId=681288] x	
	AND/OR	
) AND (		
		d O
	[	c 0
	biological process (keywor	c O
	biological process (keywor	c 0
	biological process (keywork containing c ATP binding (* support POSIX regular	c O
	( C biological process (keywork containing C ATP binding (* support POSIX regular expression) AND/OR	¢ 0
function,	( Containing Contai	0
function,	( Containing Contai	¢ 0
function,	( C biological process (keywork containing C ATP binding (* support POSIX regular expression) AND/OR ) C	¢ ♥
function,	( Containing Containining Containing Containing Containing Containing Containing Cont	0
function,	( C biological process (keywork containing C ATP binding (* support POSIX regular expression) AND/OR ) C ber starting with C 1.2.5 (* support POSIX regular	¢ 0

Supporting Figure 1. The filter box. An unlimited number of filters of different types (genomic location, presence / absence of homologs, gene ids, function, biological process, product, cellular component, EC number, type of evidence) can be combined with the operators AND (intersection) or OR (union).

## EF\_1875-EF\_1879

## EF\_2277-EF\_2281



Supporting figure 2. Example of the homolog table view showing a subset of the dispensable genes set of E. faecalis V583 and their homologs among a few compared species. The two duplicated collinear syntenies EF\_1875-EF\_1879 and EF\_2277-EF\_2281 are highlighted in red. They are conserved in our dataset among twenty-four firmicutes, all of which are pathogenic to human (see Supporting Table 4).