

Supplementary Table 1: Morbidity and Mortality Difference between included and excluded participants:

	Included in the lifetime risk analysis	
	Yes	No
n (%)	3362 (65.6)	1762 (34.4)
Age at first exam, yr (SD)	34.3 (9.4)	40.0 (11.4)
Female sex, n (%)	1765 (52.5)	876 (49.7)
CVD, n (%)	831 (24.7)	557 (31.6)
Hard CVD ² , n (%)	487 (14.5)	392 (22.2)
All-cause mortality, n (%)	599 (17.8)	949 (53.9)

¹Multivariate model adjusted for systolic blood pressure, total and HDL cholesterol, hypertension, hyperlipidemia and current smoking at exam cycle 1

²Hard CVD defined as a history of a myocardial infarction or cardiac death

Supplementary Table 2: Baseline Characteristics of Framingham Heart participants free of CKD at age 50 stratified by number of CKD risk factors

	Number of Risk Factors			
	0	1	2	3
n (%)	1793 (53.3)	1030 (30.6)	458 (13.6)	81 (2.4)
Female sex, n (%)	1028 (57.3)	500 (48.5)	198 (43.2)	39 (48.2)
Age at entry, yrs (SD)	52.3 (8.9)	57.0 (9.2)	57.7 (8.7)	59.3 (7.5)
	Baseline Characteristics			
BMI, kg/m ² (SD)	25.0 (2.8)	28.6 (4.7)	33.3 (4.9)	36.0 (4.5)
Obese, n (%)	0	376 (36.5)	385 (84.1)	81 (100)
eGFR, ml/min/1.73m ² (SD)	91.8 (15.7)	89.6 (17.8)	91.2 (18.2)	96.3 (25.5)
Hypertension, n (%)	0	608 (59.0)	412 (90.0)	81 (100)
Diabetes, n (%)	0	46 (4.5)	119 (26.0)	81 (100)
	Outcome Measures			
CKD, n (%)	267 (14.9)	281 (27.3)	152 (33.2)	29 (35.8)
Mortality, n (%)	210 (11.7)	236 (22.9)	139 (30.4)	33 (40.7)

Risk factors include the presence of obesity, hypertension or diabetes.