

Table S1. Gene-specific primer pairs used in PCR.

Gene	Primer
PIP1;3	Forward: 5'-GCTGTGGATGATCTGGTTTTATCG-3'
	Reverse: 5'-ATTTGTACCGAACCAAACCAGAAG-3'
PIP1;5	Forward: 5'-CTTCTTGTTTCCTATGTCATGTG-3'
	Reverse: 5'-GTACACAATGTATTCTTCCATTGAC-3'
PIP2;1	Forward: 5'-CAGTCACTTGCAGTTTTTCATTTCC-3'
	Reverse: 5'-GTAAACACAACGCATAAGAACCTC-3'
PIP2;3	Forward: 5'-CTTTTCCACTCGTATCTTAGCTTC-3'
	Reverse: 5'-ATACACCAAACCTTACATACGTTGC-3'
PIP2;7	Forward: 5'-TGTAATGAGAGAGATGGTGGATTG-3'
	Reverse: 5'-ACTAAGAGAAACCAAAGGCAAACG-3'

Table S2. Effects on cell elasticity (ϵ) of 5 mM $\text{Ca}(\text{NO}_3)_2$ and 5 mM KF treatments in the wild-type *Arabidopsis* plants and in plants overexpressing PIP2;5. Different letters in each column for wild-type and transgenic plants indicate significant differences (ANOVA, Turkey, $p = 0.05$). Values are means \pm SE ($n = 7$ plants).

Plants	ϵ (MPa)			
	- $\text{Ca}(\text{NO}_3)_2$	+ $\text{Ca}(\text{NO}_3)_2$	- KF	+ KF
Wild-type	4.1 ± 0.7^a	3.8 ± 0.6^a	3.7 ± 0.8^b	4.6 ± 1.2^b
Overexpressing PIP2;5-1	3.6 ± 0.9^a	3.9 ± 0.9^a	4.5 ± 0.8^b	4.6 ± 1.1^b




Figure S1. Typical responses of half-times for water exchange ($T_{1/2}$) and elastic modulus (ϵ) in cortical cells of *Arabidopsis* roots. $T_{1/2}$ is shown at about 0-20 min (A, D) and 0-35 min (B, C), and ϵ measured at about 30-40 min (A, D), 40-50 min (B, C) is shown on the right side of the traces in the control (A), KF (B), $\text{Ca}(\text{NO}_3)_2$ (C) and OA (D) treatments.




Figure S2 Confirmation of the transgenic lines. RT-PCR analyses of the expression of PIP2;5 in the wild-type (Wt) plants and independent transgenic lines (T1 and T3) in 21-day old *Arabidopsis* plants grown in hydroponic culture.