

Trace element and isotope geochemistry of Tschicoma Formation intermediate composition dome complexes, Jemez Mountains volcanic field, New Mexico, USA

Clara M. Waelkens, John Stix, Fraser Goff & Dominique Weis

Supplementary table: Overview of ages of the Tschicoma lava dome and lava flow complexes and locations where they are preserved.

Lava dome complex	Location	Age
Tschicoma Peak area	North and northeast of caldera	2.62 ± 0.64 Ma – 5.34 ± 0.36 Ma (Kemper <i>et al.</i> , 2007)
Caballo Mountain	Sierra de los Valles	3.06 ± 0.15 Ma – 4.66 ± 0.17 Ma (Broxton <i>et al.</i> , 2007)
Cerro Rubio	Toledo embayment	3.56 ± 0.36 Ma – 3.59 ± 0.36 Ma (K-Ar ages) (Heiken <i>et al.</i> , 1986; Stix <i>et al.</i> , 1988)
Upper Quemazon	Eastern caldera rim	2.92 ± 0.03 Ma (Kelley <i>et al.</i> , 2013)
Cañada Bonita	Eastern caldera rim	3.50 ± 0.23 Ma (Kelley <i>et al.</i> , 2013; Goff <i>et al.</i> , 2021)
Rendija Canyon	Sierra de los Valles	5.038 ± 0.003 Ma – 5.062 ± 0.006 Ma (Goff <i>et al.</i> , 2021)
Pajarito Mountain	Sierra de los Valles	2.93 ± 0.06 Ma – 3.09 ± 0.08 Ma (Broxton <i>et al.</i> , 2007)
Cerro Grande	Sierra de los Valles	2.88 ± 0.02 Ma – 3.35 ± 0.17 Ma (Broxton <i>et al.</i> , 2007)
Sawyer Dome	Sierra de los Valles	3.18 ± 0.20 Ma – 3.67 ± 0.29 Ma (Goff <i>et al.</i> , 2006; Broxton <i>et al.</i> , 2007; Kelley <i>et al.</i> , 2013)
Redondo Peak megabreccia	Redondo Peak resurgent dome	2.01 ± 0.02 Ma – 2.20 ± 0.05 Ma (Kelley <i>et al.</i> , 2013)